



Orni Super Mix **Extra Strong**

Ornithosis, Mycoplasmosis, Respiratory problems and Canker.

Symptoms

Picking of feathers, scratching head, red throat, yawning, inflamed third eyelid (especially in young pigeons), dirty noses, blue flesh, swollen heads, one eye cold, wet eyes, snot.

Dosage

Prevention: 1 measuring spoon (=5g) for 2 litres drinking water during 1 or 2 days every 2 weeks, preferably starting the day after the race.

As a treatment: 1 measuring spoon (=5g) for 2 litres drinking water during 5-7 successive days. This cure may be repeated if necessary after a few days.

Contains: Doxycycline Hyclate 6%, Spiramycin Adipate 10%, Metronidazole 10%, Bromhexine HCl 2%, Inactive ingredients q.s. to 100 g.

100 g POWDER

50 pcs TABS

Dosage

Preventive: 1 tablet per pigeon (400-500 g bodyweight) for 1-2 consecutive days.

Curative: 1 tablet per pigeon (400-500 g bodyweight) for 3-5 consecutive days, or as vet. prescribes.

Contains: Doxycycline Hyclate 102 mg/g ; Spiramycin Adipate 80 mg/g ; Metronidazole 100 mg/g ; Bromhexine HCl 20 mg/g